

COMPARISON BETWEEN TEICHMÜLLER AND LIPSCHITZ METRICS

YOUNG-EUN CHOI AND KASRA RAFI

1. INTRODUCTION

The Teichmüller distance between two points σ, τ in Teichmüller space $\mathcal{T}(S)$ is defined in terms of the minimal quasiconformal constant $K(\sigma, \tau)$ between σ and τ . In [10] Thurston introduced an analogous metric on $\mathcal{T}(S)$ by considering the least possible value of the global Lipschitz constant $\Lambda(\sigma, \tau)$ from σ to τ . On the one hand, Kerckhoff [3] showed that $K(\sigma, \tau)$ can be formulated in terms of the ratio of extremal lengths of simple closed curves:

$$(1) \quad K(\sigma, \tau) = \sup_{\alpha} \frac{\text{Ext}_{\tau}(\alpha)}{\text{Ext}_{\sigma}(\alpha)}$$

and on the other, it was shown by Thurston [10] that the minimal Lipschitz constant $\Lambda(\sigma, \tau)$ is given by the ratio of lengths in the hyperbolic metric:

$$(2) \quad \Lambda(\sigma, \tau) = \sup_{\alpha} \frac{l_{\tau}(\alpha)}{l_{\sigma}(\alpha)}.$$

A comparison of $K(\sigma, \tau)$ and the ratio of lengths in Equation (2) was first given by Wolpert [11], who proved that for any K -quasiconformal map f from σ to τ and any simple closed curve α

$$\frac{l_{\tau}(f(\alpha))}{l_{\sigma}(\alpha)} \leq K.$$

This implies, in particular, that

$$\Lambda(\sigma, \tau) \leq K(\sigma, \tau).$$

In this paper, we compare the Teichmüller and Lipschitz metrics by comparing the two ratios in Equations (1) and (2). Our method is to analyze the ratio of hyperbolic lengths in much the same way the ratio of extremal lengths was analyzed by Minsky in [5] for the purpose of showing that certain regions in the thin part of Teichmüller space have product structures. However, since $K(\sigma, \tau)$ is symmetric and $\Lambda(\sigma, \tau)$ is

not [10], it is necessary to choose some symmetric version of Λ to make the comparison more meaningful. Thus, we take

$$L(\sigma, \tau) = \max\{\Lambda(\sigma, \tau), \Lambda(\tau, \sigma)\}$$

and define the Teichmüller and Lipschitz metrics, respectively, as follows

$$\begin{aligned} d_{\mathcal{T}}(\sigma, \tau) &= \frac{1}{2} \log K(\sigma, \tau) \\ d_L(\sigma, \tau) &= \log L(\sigma, \tau). \end{aligned}$$

Note that the factor of $1/2$ has been left out in the Lipschitz metric. This is due to the fact that on the thick part of Teichmüller space, we can compare the two metrics up to an additive error, as we shall shortly see.

Although $\Lambda(\sigma, \tau)$ is not symmetric, it is easy to check that it satisfies the following ordered triangle inequality:

$$\Lambda(\rho, \tau) \leq \Lambda(\rho, \sigma) + \Lambda(\sigma, \tau)$$

and further, satisfies the property that $\Lambda(\sigma, \tau) = 0$ if and only if $\sigma = \tau$. Thus $d_L(\sigma, \tau)$ defines a genuine metric, in that it is symmetric, takes the value zero if and only if $\sigma = \tau$, and satisfies the triangle inequality. In [1], it was shown that on the Teichmüller space of the torus, the Teichmüller metric and a similarly defined Lipschitz metric are, in fact, equal. In contrast, we show that for a hyperbolic surface S , the two metrics are not comparable. In particular,

Theorem A. *There are sequences $\sigma_n, \tau_n \in \mathcal{T}(S)$ such that, as $n \rightarrow \infty$,*

$$d_L(\sigma_n, \tau_n) \rightarrow 0, \quad d_{\mathcal{T}}(\sigma_n, \tau_n) \rightarrow \infty.$$

As is often the case, however, no incongruities occur on the thick part of Teichmüller space, and there the two metrics are quasi-isometric to one another. In fact, they are equal up to a bounded additive error. This is a consequence of the following theorem proved in Section 2:

Theorem B. *For $\rho \in \mathcal{T}(S)$, let μ_ρ be a short marking for ρ . For every $\epsilon > 0$, there is a constant c depending on ϵ such that, for any σ, τ in the ϵ -thick part of $\mathcal{T}(S)$, the following quantities differ from one another by at most c :*

- | | |
|--|--|
| 1. $d_{\mathcal{T}}(\sigma, \tau)$ | 2. $d_L(\sigma, \tau)$ |
| 3. $\log \max_{\alpha \in \mu_\sigma} \frac{l_\tau(\alpha)}{l_\sigma(\alpha)}$ | 4. $\log \max_{\alpha \in \mu_\tau} \frac{l_\sigma(\alpha)}{l_\tau(\alpha)}$. |

In particular, in order to estimate the Teichmüller distance between two points in the thick part, one need only compare the lengths of a finite number of curves with respect to the two metrics.

To compare the metrics on the thin part of Teichmüller space, we prove in Section 3 an analogue of Minsky's product region theorem [5]. Let Γ be a collection of k disjoint, homotopically distinct, simple closed curves on S and let $Thin_\epsilon(S, \Gamma)$ be the set of $\sigma \in \mathcal{T}(S)$ such that $l_\sigma(\gamma) \leq \epsilon$ for all $\gamma \in \Gamma$. Let $T_\Gamma = \mathcal{T}(S \setminus \Gamma) \times U_1 \times \cdots \times U_k$, where $S \setminus \Gamma$ is the analytically finite surface obtained from S by pinching all the curves in Γ and where U_i is the subset $\{(x, y) : y \geq 1/\epsilon\}$ of the upper-half plane. The Fenchel-Nielsen coordinates on $\mathcal{T}(S)$ give rise to a natural homeomorphism $\Pi : Thin_\epsilon(S, \Gamma) \rightarrow T_\Gamma$. Then Minsky's product region theorem states:

Theorem 1.1 (Minsky [5]). *Let d_{T_Γ} be the sup metric*

$$d_{T_\Gamma} = \sup \left\{ d_{\mathcal{T}(S \setminus \Gamma)}, \frac{1}{2}d_{\mathbb{H}_1}, \dots, \frac{1}{2}d_{\mathbb{H}_k} \right\}$$

on T_Γ , where $d_{\mathcal{T}(S \setminus \Gamma)}$ is the Teichmüller metric on $\mathcal{T}(S \setminus \Gamma)$ and $d_{\mathbb{H}_i}$ is the restriction of the hyperbolic metric on the upper-half plane to U_i . Then, for ϵ sufficiently small, there is a constant c depending on ϵ , such that for any $\sigma, \tau \in Thin_\epsilon(S, \Gamma)$,

$$|d_{\mathcal{T}}(\sigma, \tau) - d_{T_\Gamma}(\Pi(\sigma), \Pi(\tau))| < c.$$

In the analogue for the Lipschitz metric, we define the sup metric

$$d_{L_\Gamma} = \sup \{d_{L(S \setminus \Gamma)}, d_{L(A_1)}, \dots, d_{L(A_k)}\}$$

on T_Γ , where $d_{L(S \setminus \Gamma)}$ is the Lipschitz metric on $\mathcal{T}(S \setminus \Gamma)$ and $d_{L(A_i)}$ is a modification of the hyperbolic metric on U_i (see Section 3 for details):

Theorem C. *For ϵ sufficiently small, there is a constant c depending on ϵ , such that for any $\sigma, \tau \in Thin_\epsilon(S, \Gamma)$,*

$$|d_L(\sigma, \tau) - d_{L_\Gamma}(\Pi(\sigma), \Pi(\tau))| < c.$$

A more precise statement is given in Theorem 3.4. Our proof follows parallel to Minsky's, but requires only elementary hyperbolic geometry, since we need not deal with extremal lengths.

As a consequence of Theorem B, one can deduce the following purely combinatorial result. For a subsurface Z , let $d_Z(\mu_1, \mu_2)$ be the distance between the projections of μ_1 and μ_2 to Z , measured in the arc complex of Z (see [4], [8] for details).

Corollary D. *There is a constant k such that for any markings μ_1 and μ_2 on S ,*

$$(3) \quad \log i(\mu_1, \mu_2) \asymp \sum_Y [d_Y(\mu_1, \mu_2)]_k + \sum_A \log [d_A(\mu_1, \mu_2)]_k,$$

where Y ranges over all subsurfaces of S that are not annuli, A ranges over all annuli, and where $[x]_k = 0$ if $x < k$ and $[x]_k = x$ if $x \geq k$.

In [4], Masur and Minsky provide an estimate, similar to the right-hand side of (3), for the number of elementary moves needed to change μ_1 to μ_2 . Using their result and examining how the intersection number between two markings changes as a result of applying a sequence of elementary moves to one of them, one can show that the right-hand side of (3) is an upper bound for $\log i(\mu_1, \mu_2)$ (there is no clear combinatorial argument for proving the inequality in the other direction). In this context, Corollary D states that, along an efficient path in the marking space, the intersection number increases at the fastest possible rate.

1.1. Notation. Often, we shall compare two functions f, g on $\mathcal{T}(S)$ and use the notation $f \prec g$, $f \asymp g$ to mean, respectively, that there are positive constants k, c such that $f \leq kg + c$, $\frac{1}{k}g - c \leq f \leq kg + c$. We also use $f \dot{\prec} g$, $f \dot{\asymp} g$ to mean, respectively, that there is only a multiplicative constant, or only an additive constant, involved. In particular, $f \dot{\asymp} 1$ means that the function f is bounded both above and below by positive constants. The constants k and c usually depend on the topological type of S , which will not be subsequently mentioned. Any other dependencies will be explicitly noted.

2. THE THICK PART

Let S be a surface of finite topological type. Given $\epsilon > 0$, the ϵ -thick part of Teichmüller space is the set of $\sigma \in \mathcal{T}(S)$ such that the infimum of the injectivity radius measured in σ , taken over all points in S , is greater than ϵ . When we simply say “the thick part”, we mean it is the ϵ -thick part for some ϵ which has already been chosen.

A *marking* on S is a collection of homotopically distinct, simple closed curves in S obtained by first choosing a pants curves system, i.e., a collection of mutually disjoint curves that cut S into pairs of pants (where a hole may be a puncture of S) and then by choosing an additional collection of curves that together with the pants system cuts the surface into disks and punctured disks. To make the choice of a marking less arbitrary, additional conditions on the choice of curves are often specified.

For $\sigma \in \mathcal{T}(S)$, we define a *short marking* μ_σ , as follows. First choose a pants system by taking the shortest curve in S , then the next shortest curve disjoint from the first, and so on until a complete pants system $\underline{\alpha}$ is formed. We remark that throughout this paper, when we say the “length of a curve”, we always mean the length of its geodesic representative. Next, choose a “dual” curve δ_α for each $\alpha \in \underline{\alpha}$ that is disjoint from $\underline{\alpha} \setminus \alpha$, and that is shortest among all such curves. There may be a finite number of possible short markings for σ .

A lemma of Bers’ says that there is a uniform constant N such that every $\sigma \in \mathcal{T}(S)$ has a pants curves system $\underline{\alpha}$ with the property that $l_\sigma(\alpha) < N$ for all $\alpha \in \underline{\alpha}$. Hence, if σ is in the ϵ -thick part of $\mathcal{T}(S)$ so that all the curves in a short marking μ have length bounded below as well, then the lengths of the dual curves are bounded above and so $l_\sigma(\mu)$ is bounded above by some quantity depending only on ϵ . Conversely, given a marking μ and a number $B > 0$, the metrics $\sigma \in \mathcal{T}(S)$ such that $l_\sigma(\mu) = \sum_{\alpha \in \mu} l_\sigma(\alpha) \leq B$ has bounded diameter in $\mathcal{T}(S)$, where the bound depends only on B (see for example [6]). Thus there is a coarse correspondence between the thick part of Teichmüller space and the set of markings. This idea is implicit in the theorems to follow.

Theorem B. *For every $\epsilon > 0$, there is a constant c depending on ϵ such that, for any σ, τ in the ϵ -thick part of $\mathcal{T}(S)$, the following quantities differ from one another by at most c :*

1. $d_{\mathcal{T}}(\sigma, \tau)$
2. $d_L(\sigma, \tau)$
3. $\log \max_{\alpha \in \mu_\sigma} \frac{l_\tau(\alpha)}{l_\sigma(\alpha)}$
4. $\log \max_{\alpha \in \mu_\tau} \frac{l_\sigma(\alpha)}{l_\tau(\alpha)}$.

First we need the following lemma. Let $g : \mathbb{R} \rightarrow \mathcal{T}(S)$ be the Teichmüller geodesic that passes through σ and τ and let q_t be the family of quadratic differentials representing g . We assume all quadratic differential metrics have been normalized to have area 1.

Lemma 2.1. *For every marking μ on S there exist l_0 and t_0 such that*

$$l_{q_t}(\mu) \asymp l_0 e^{|t-t_0|}.$$

Proof. Recall that a quadratic differential q_t defines a pair of measured foliations on the surface S , called the horizontal and the vertical foliations. For every curve α the horizontal length $h_t(\alpha)$ of α is the intersection number of α with the horizontal foliation and the vertical length $v_t(\alpha)$ of α is the intersection number of α with the vertical foliation. We have (see for example [7])

$$l_{q_t}(\alpha) \asymp h_t(\alpha) + v_t(\alpha).$$

Let t_α be the time when α is balanced, i.e., the time when the horizontal length and the vertical length of α are equal. Let $l_\alpha = l_{q_{t_\alpha}}(\alpha)$. Along a Teichmüller geodesic, the horizontal length of α increases and the vertical length of α decreases exponentially fast. Therefore,

$$l_{q_t}(\alpha) \asymp l_\alpha \cosh(t - t_\alpha).$$

Thus, for every marking μ

$$(4) \quad l_{q_t}(\mu) = \sum_{\alpha \in \mu} l_{q_t}(\alpha) \asymp \sum_{\alpha \in \mu} l_\alpha \cosh(t - t_\alpha).$$

Denote the right hand side of (4) by $f(t)$. Let t_0 be the time when $f(t)$ is minimum and let $l_0 = f(t_0)$. Since

$$\cosh(t - t_\alpha) \leq \cosh(t_0 - t_\alpha) e^{|t-t_0|},$$

we have

$$(5) \quad \sum_{\alpha \in \mu} l_\alpha \cosh(t - t_\alpha) \leq \sum_{\alpha \in \mu} l_\alpha \cosh(t_0 - t_\alpha) e^{|t-t_0|} = l_0 e^{|t-t_0|}.$$

To prove the inequality in the other direction, we observe that the derivative of $f(t)$ with respect to t at $t = t_0$ is $\sum_{\alpha} l_\alpha \sinh(t_0 - t_\alpha) = 0$, which implies

$$\sum_{\alpha \in \mu} l_\alpha e^{t_0 - t_\alpha} = \sum_{\alpha \in \mu} l_\alpha e^{t_\alpha - t_0} = l_0.$$

If n is the number of curves in μ , the above equation implies that there exist $\beta, \gamma \in \mu$ such that

$$l_\beta e^{t_0 - t_\beta} \geq \frac{l_0}{n} \quad \text{and} \quad l_\gamma e^{t_\gamma - t_0} \geq \frac{l_0}{n}.$$

Thus we have

$$\begin{aligned} f(t) &= \sum_{\alpha \in \mu} l_\alpha \cosh(t - t_\alpha) \geq l_\beta \cosh(t - t_\beta) + l_\gamma \cosh(t - t_\gamma) \\ &\geq \frac{1}{2} \left[l_\beta e^{t-t_0} e^{t_0-t_\beta} + l_\gamma e^{t_\gamma-t_0} e^{t_0-t} \right] \\ (6) \quad &\geq \frac{l_0}{2n} e^{|t-t_0|}. \end{aligned}$$

Equations (5) and (6) show that $f(t) \asymp l_0 e^{|t-t_0|}$. This and (4) prove the lemma. $\square$

Proof of Theorem B. We show that the first three quantities are comparable, the proof for the remaining term is similar. Suppose that for $a < b$, we have $g(a) = \sigma$, $g(b) = \tau$ so that $d_{\mathcal{T}}(\sigma, \tau) = b - a$. Since the moduli space of the thick part is compact, we know that the hyperbolic lengths of curves in σ, τ are proportional to their quadratic differential

lengths in q_a, q_b , respectively (see [9] for a more general discussion). Therefore, there are multiplicative constants depending only on ϵ , such that for any simple closed curve α ,

$$(7) \quad \frac{l_\tau(\alpha)}{l_\sigma(\alpha)} \asymp \frac{l_{q_b}(\alpha)}{l_{q_a}(\alpha)}.$$

Moreover, since

$$l_{q_b}(\alpha) \asymp l_\alpha \cosh(b - t_\alpha) \leq e^{b-a} l_\alpha \cosh(a - t_\alpha) = e^{b-a} l_{q_a}(\alpha),$$

it follows from Equation (7) that

$$d_L(\sigma, \tau) \stackrel{+}{\prec} d_T(\sigma, \tau).$$

Thus it remains to be shown that there is a curve $\alpha \in \mu_\sigma$ such that

$$b - a \stackrel{+}{\prec} \log \frac{l_{q_b}(\alpha)}{l_{q_a}(\alpha)}.$$

Let $l_{q_t}(\mu_\sigma) \asymp l_0 e^{|t-t_0|}$ as in Lemma 2.1. Then,

$$(8) \quad l_0 e^{|b-a|-|a-t_0|} \prec l_{q_b}(\mu_\sigma) \prec l_0 e^{|b-a|+|a-t_0|}.$$

First we show that $|a-t_0|$ is bounded above. Since σ is in the thick part, the q_a -length and the σ -length of μ_σ are comparable to one another. Moreover, since μ_σ is a short marking in σ , its σ -length is bounded both above and below. Therefore, we have:

$$(9) \quad l_0 e^{|a-t_0|} \asymp l_{q_a}(\mu_\sigma) \asymp l_\sigma(\mu_\sigma) \asymp 1.$$

Furthermore, we can see that l_0 is bounded below, as follows. A marking divides the surface into disks and punctured disks. For any quadratic differential q , the q -area of a disk or a punctured disk is less than the square of its perimeter. Therefore, we have for all t ,

$$(10) \quad 1 = \text{area}_{q_t}(S) \prec \sum_{\alpha \in \mu_\sigma} l_{q_t}(\alpha)^2.$$

Applied to $t = t_0$ we get $l_0 \succ 1$. It then follows from Equation (9) that $|a - t_0| \prec 1$, as desired. Thus, it follows from Equation (8) that

$$l_{q_b}(\mu_\sigma) \asymp e^{b-a}.$$

But, as we saw in Equation (9), since the q_a -lengths of curves in μ_σ are bounded above and below, it follows that there exists a curve $\alpha \in \mu_\sigma$ such that

$$l_{q_b}(\alpha) \asymp l_{q_a}(\alpha) e^{b-a},$$

which is what we wanted. $\square$

Theorem 2.2. *Let σ and τ be points in the ϵ -thick part of Teichmüller space and let μ_σ and μ_τ be their short markings, respectively. Then there is an additive constant depending only on ϵ , such that*

$$d_{\mathcal{T}}(\sigma, \tau) \stackrel{+}{\asymp} \log i(\mu_\sigma, \mu_\tau),$$

where $i(\mu_\sigma, \mu_\tau)$ is the total number of intersections between the curves in μ_σ and the curves in μ_τ .

Proof. The τ -length of a curve is proportional to its intersection number with μ_τ (see for example [6, Lemma 4.7]). Therefore,

$$(11) \quad i(\mu_\sigma, \mu_\tau) \asymp \sum_{\alpha \in \mu_\sigma} l_\tau(\alpha) \asymp \max_{\alpha \in \mu_\sigma} l_\tau(\alpha).$$

Since σ is in the thick part of $\mathcal{T}(S)$, we have $l_\sigma(\alpha) \asymp 1$ for every curve $\alpha \in \mu_\sigma$. Thus, it follows from Theorem B that

$$(12) \quad \log \max_{\alpha \in \mu_\sigma} l_\tau(\alpha) \stackrel{+}{\asymp} \log \max_{\alpha \in \mu_\sigma} \frac{l_\tau(\alpha)}{l_\sigma(\alpha)} \stackrel{+}{\asymp} d_{\mathcal{T}}(\sigma, \tau).$$

The theorem follows from Equations (11) and (12). $\square$

Remark 2.3. *The above theorem implies that the logarithm of the intersection number is almost a distance function on the marking space. In particular, it satisfies a quasi-triangle inequality. That is, for markings μ_1 , μ_2 and μ_3 we have*

$$\log i(\mu_1, \mu_3) \stackrel{+}{\preceq} \log i(\mu_1, \mu_2) + \log i(\mu_2, \mu_3).$$

This “distance function” is similar, but not comparable to the distance defined on the space of markings in [4].

Proof of Corollary D. For given markings μ_1 and μ_2 , one can find points σ_1 and σ_2 in the thick part of Teichmüller space such that μ_1 and μ_2 are short markings in σ_1 and σ_2 , respectively. In [8], a combinatorial formula is given for the Teichmüller distance between two points in the thick part of Teichmüller space. It states that $d_{\mathcal{T}}(\sigma_1, \sigma_2)$ is comparable to the right-hand side of Equation (3). Also, Theorem 2.2 states that $\log i(\mu_1, \mu_2) \stackrel{+}{\asymp} d_{\mathcal{T}}(\sigma_1, \sigma_2)$. Together these two results prove the corollary. $\square$

3. PRODUCT REGIONS IN THE LIPSCHITZ METRIC

In this section, we prove the analogue of Minsky’s product region theorem for the Lipschitz metric.

3.1. An (ϵ_0, ϵ_1) -decomposition. First, we need to recall the notion of an (ϵ_0, ϵ_1) -decomposition defined in [5]. Let $0 < \epsilon_1 < \epsilon_0$ be two numbers less than the Margulis constant. Let σ be a hyperbolic metric on S and suppose $\gamma_1, \dots, \gamma_k$ are geodesics with length $l_\sigma(\gamma_i) \leq \epsilon_1$. Let $A_1, \dots, A_k$ be the collection of annular neighborhoods of $\gamma_1, \dots, \gamma_k$, respectively, such that the boundary components of A_i each have length ϵ_0 . A component Q of $S \setminus \cup A_i$ is called a *hyperbolic* component and the entire collection $\mathcal{P}$ of hyperbolic components and annular components is called an (ϵ_0, ϵ_1) -decomposition. We assume that ϵ_0, ϵ_1 are chosen so that any simple geodesic that intersects an annular component A is either the core of A or is made up of arcs that run from one boundary component of A to another. We remark that in [5], what we have described is called a *partial* (ϵ_0, ϵ_1) -decomposition. There, the term (ϵ_0, ϵ_1) -decomposition is reserved for the case where $\{\gamma_1, \dots, \gamma_k\}$ is the full set of curves whose length l_σ satisfies $l_\sigma \leq \epsilon_1$.

In the course of arguments to follow, we shall further require that $\epsilon_0/\epsilon_1 > 2$ so that certain desired estimates hold (see for example Lemma 3.5). We therefore assume ϵ_0, ϵ_1 have been chosen once and for all to satisfy all the conditions stated above and henceforth use the notation $f \asymp g$, $f \stackrel{+}{\asymp} g$, etc., to mean that the multiplicative or additive constants which appear depend only on this choice of ϵ_0, ϵ_1 (and on the topological type of S).

3.2. Decomposing the length of a curve. Consider the intersection of a simple closed curve ζ with the components of an (ϵ_0, ϵ_1) -decomposition. For a hyperbolic component Q , let $\mathcal{C}(Q, \partial Q)$ denote the homotopy classes of simple closed curves in Q and of essential arcs in Q with endpoints on ∂Q , under homotopies that keep any endpoints of arcs on ∂Q . Define the orthogonal projection ζ_Q of ζ to be the geodesic representative of $\zeta \cap Q$ in $\mathcal{C}(Q, \partial Q)$ that has the shortest length (see [5, §2.3]). In particular, every arc in ζ_Q is perpendicular to ∂Q . It is not hard to show the following:

Proposition 3.1. *Let $\mathcal{P}$ be the components of an (ϵ_0, ϵ_1) -decomposition for σ and let $Q, A \in \mathcal{P}$ be respectively, a hyperbolic and annular component. Then, for any simple closed curve ζ , the following estimates hold:*

$$(13) \quad i(\zeta, \partial Q) \asymp |l_\sigma(\zeta \cap Q) - l_\sigma(\zeta_Q)|$$

$$(14) \quad i(\zeta, \gamma) \asymp \left| l_\sigma(\zeta \cap A) - \left[\log \frac{\epsilon_0}{l_\sigma(\gamma)} + l_\sigma(\gamma) \frac{Tw_\sigma(\zeta, \gamma)}{2} \right] i(\zeta, \gamma) \right|$$

where γ is the core geodesic of A .

Here, $Tw_\sigma(\zeta, \gamma)$ is the absolute value of the twist of ζ around γ defined in [5, §3]. In Equation (14), $\log[\epsilon_0/l_\sigma(\gamma)]$ is approximately half the width of A ; the right-hand side describes the sum of lengths of piecewise geodesic arcs homotopic to $\zeta \cap A$ in $\mathcal{C}(A, \partial A)$, each of which goes perpendicularly from one component of A to γ , wraps around γ a number of $Tw_\sigma(\zeta, \gamma)$ times (up to an error of 1), then goes out the other end of A orthogonally. The idea is that most of the twisting ζ does around γ , takes place in A [5]. This is also the reason that Equation (13) is true. For a proof, see [2].

Since the components of ∂Q each have a collar of some definite width, $l_\sigma(\zeta \cap Q) \asymp i(\zeta, \partial Q)$ and $l_\sigma(\zeta_Q) \asymp i(\zeta, \partial Q)$. Similarly, since γ has a collar of definite width, terms in the righthand side of Equation (14) are larger than a multiple of $i(\zeta, \gamma)$. Therefore, we can rewrite Equations (13) and (14) as follows:

Corollary 3.2. *Let Q, A be as in Proposition 3.1. Then for any simple closed curve ζ on S , we have*

$$(15) \quad l_\sigma(\zeta \cap Q) \asymp l_\sigma(\zeta_Q)$$

$$(16) \quad l_\sigma(\zeta \cap A) \asymp \left[\log \frac{\epsilon_0}{l_\sigma(\gamma)} + l_\sigma(\gamma) \cdot \frac{Tw_\sigma(\zeta, \gamma)}{2} \right] i(\zeta, \gamma).$$

3.3. Regular Annuli. Let A be an annulus. We call a metric ρ on A a *regular metric* if (A, ρ) is isometric to quotient of some closed neighborhood $\{ p \in \mathbb{H}^2 \mid d_{\mathbb{H}^2}(p, G) \leq r \}$ of a geodesic G in $\mathbb{H}^2$, by a hyperbolic isometry with axis G . For $\epsilon > 0$, let $U_\epsilon(A)$ be the space of all regular metrics on A such that the core of A has length at most ϵ and such that each component of ∂A has length ϵ_0 . Two metrics are considered equivalent if they differ by an isotopy of A fixing ∂A pointwise. Define the distance between $\rho_1, \rho_2 \in U_\epsilon(A)$ to be

$$d_{L(A)}(\rho_1, \rho_2) = \sup_{\beta \in \mathcal{C}(A, \partial A)} \left| \log \frac{l_{\rho_1}(\beta)}{l_{\rho_2}(\beta)} \right|,$$

where $\mathcal{C}(A, \partial A)$ is the set of homotopy classes of non-trivial simple loops or arcs in A , under homotopies that fix the endpoints. As usual, the length $l_\rho(\beta)$ means the length of the ρ -geodesic representative of β . Clearly $d_{L(A)}(\rho_1, \rho_2)$ is symmetric, and is zero if and only if $\rho_1 = \rho_2$.

To see that the triangle inequality holds, observe that

$$\begin{aligned} \left| \log \frac{l_{\rho_1}(\beta)}{l_{\rho_2}(\beta)} \right| + \left| \log \frac{l_{\rho_2}(\beta)}{l_{\rho_3}(\beta)} \right| &\geq \log \frac{l_{\rho_1}(\beta)}{l_{\rho_2}(\beta)} + \log \frac{l_{\rho_2}(\beta)}{l_{\rho_3}(\beta)} = \log \frac{l_{\rho_1}(\beta)}{l_{\rho_3}(\beta)}, \\ \left| \log \frac{l_{\rho_1}(\beta)}{l_{\rho_2}(\beta)} \right| + \left| \log \frac{l_{\rho_2}(\beta)}{l_{\rho_3}(\beta)} \right| &\geq \log \frac{l_{\rho_2}(\beta)}{l_{\rho_1}(\beta)} + \log \frac{l_{\rho_3}(\beta)}{l_{\rho_2}(\beta)} = \log \frac{l_{\rho_3}(\beta)}{l_{\rho_1}(\beta)}. \end{aligned}$$

Let γ be the core of A and fix a simple arc ω in A that connects the two components of ∂A . For every $\rho \in U_\epsilon(A)$, the twist parameter $tw_\rho(A)$ of ρ is defined as follows (see also [5, §3]). First, it is necessary to fix an orientation of γ . Consider the universal cover of (A, ρ) in $\mathbb{H}^2$ and the lifts $\tilde{\gamma}$, $\tilde{\omega}$ of γ, ω , respectively (see Figure 1). Extend $\tilde{\omega}$




FIGURE 1. Defining twist parameter.

to an infinite geodesic $\bar{\omega}$ and let ω_L, ω_R be the endpoints of $\bar{\omega}$ that lie on the left and right of $\tilde{\gamma}$, respectively. Let p_L, p_R be respectively, the orthogonal projections of ω_L, ω_R to $\tilde{\gamma}$. Then the twist parameter is defined as

$$tw_\rho(A) = \pm \frac{d_{\mathbb{H}^2}(p_L, p_R)}{l_\rho(\gamma)},$$

where the sign is $(+)$ if the direction from p_L to p_R coincides with the orientation of $\tilde{\gamma}$ and $(-)$ if it is opposite.

Then $U_\epsilon(A)$ can be parameterized by the length of γ and the twist parameter. The map $\rho \mapsto (tw_\rho(A), 1/l_\rho(\gamma))$ is a homeomorphism identifying $U_\epsilon(A)$ with a subset of the upper half plane:

$$U_\epsilon(A) = \left\{ (x, y) \in \mathbb{R}^2 \mid y \geq \frac{1}{\epsilon} \right\}.$$

We can formulate the distance $d_{L(A)}$ on $U_\epsilon(A)$ in terms of these coordinates as follows. Let $\rho_1, \rho_2 \in U_\epsilon(A)$ and let $t_i = tw_{\rho_i}(A)$, $l_i = l_{\rho_i}(\gamma)$ for $i = 1, 2$.

Lemma 3.3. *Assume that $l_1 \leq l_2$. Then the following hold.*

(i) If $|t_1 - t_2| l_1 \leq \log[1/l_1]$, then

$$d_{L(A)}(\rho_1, \rho_2) \stackrel{+}{\asymp} \log \frac{l_2}{l_1}.$$

(ii) If $|t_1 - t_2| l_1 > \log[1/l_1]$, then

$$d_{L(A)}(\rho_1, \rho_2) \stackrel{+}{\asymp} \log \frac{|t_1 - t_2| l_2}{\log[1/l_1]} = \log \frac{l_2}{l_1} + \log \frac{|t_1 - t_2| l_1}{\log[1/l_1]}.$$

We remark that in comparison, the hyperbolic distance between $z_1 = (t_1, 1/l_1)$ and $z_2 = (t_2, 1/l_2)$ in the upper-half plane can be estimated as follows. Assume that $l_1 \leq l_2$.

(i) If $|t_1 - t_2| l_1 \leq 1$ then

$$d_{\mathbb{H}^2}(z_1, z_2) \stackrel{+}{\asymp} \log \frac{l_2}{l_1}.$$

(ii) If $|t_1 - t_2| l_1 > 1$ then

$$d_{\mathbb{H}^2}(z_1, z_2) \stackrel{+}{\asymp} \log \frac{l_2}{l_1} + 2 \log[|t_1 - t_2| l_1].$$

Proof. For any arc $\beta \in \mathcal{C}(A, \partial A)$ intersecting γ and for $\rho \in U_\epsilon(A)$, we can define the twist $tw_\rho(\beta, \gamma)$ of β around the (oriented) curve γ in the same way we defined $tw_\rho(A)$, by replacing the reference arc ω with β in that construction. It follows from Corollary 3.2 that

$$\frac{l_2(\beta)}{l_1(\beta)} \stackrel{\cdot}{\asymp} \frac{\log[1/l_2] + |tw_{\rho_2}(\beta, \gamma)| l_2}{\log[1/l_1] + |tw_{\rho_1}(\beta, \gamma)| l_1}.$$

Moreover, it follows from [5, Lemma 3.5] that

$$|[tw_{\rho_2}(\beta, \gamma) - tw_{\rho_1}(\beta, \gamma)] - [t_2 - t_1]| \stackrel{+}{\asymp} 1.$$

Thus, the supremum over all arcs β intersecting γ is

$$(17) \quad \sup_{\beta \in \mathcal{C}(A, \partial A)} \frac{l_2(\beta)}{l_1(\beta)} \stackrel{\cdot}{\asymp} \max \left\{ \frac{l_2}{l_1}, \frac{\log[1/l_2] + |t_2 - t_1| l_2}{\log[1/l_1]} \right\}.$$

To simplify notation, let

$$R_1 = \frac{\log[1/l_2] + |t_2 - t_1| l_2}{\log[1/l_1]}, \quad R_2 = \frac{\log[1/l_1] + |t_2 - t_1| l_1}{\log[1/l_2]},$$

and

$$R = \frac{|t_2 - t_1| l_2}{\log[1/l_1]}.$$

The assumption that $l_1 \leq l_2$ implies that $R_1 \asymp R$ and that $R_2 < l_2/l_1 + R$. Therefore,

$$d_{L(A)}(\rho_1, \rho_2) \stackrel{+}{\asymp} \log \max \left\{ R, \frac{l_2}{l_1} \right\}.$$

If $|t_1 - t_2| l_1 \leq \log[1/l_1]$, then $R < l_2/l_1 + 1$ so

$$d_{L(A)}(\rho_1, \rho_2) \stackrel{+}{\asymp} \log l_2/l_1.$$

If $|t_1 - t_2| l_1 > \log[1/l_1]$, then $R > l_2/l_1$ and hence

$$d_{L(A)}(\rho_1, \rho_2) \stackrel{+}{\asymp} \log R. \quad \square$$

3.4. Statement of theorem. Let $\Gamma = \{\gamma_1, \dots, \gamma_k\}$ be a collection of disjoint, homotopically distinct simple closed curves on S and let $A_1, \dots, A_k$ be collars around $\gamma_1, \dots, \gamma_k$, respectively. Choose a Fenchel-Nielsen coordinate system associated to a marking that contains Γ in its pants system. Let $s_\sigma(\gamma_i)$ denote the Fenchel-Nielsen twist coordinate of γ_i . Let $U_i = U_{\epsilon_1}(A_i)$. For $\sigma \in \text{Thin}_{\epsilon_1}(S, \Gamma)$, let $\Pi_i(\sigma) \in U_i$ be the metric ρ whose twist $tw_\rho(A) = s_\sigma(\gamma_i)$ and such that $l_\rho(\gamma_i) = l_\sigma(\gamma_i)$. Each $\sigma \in \text{Thin}_{\epsilon_1}(S, \Gamma)$ also defines a metric $\Pi_{S \setminus \Gamma}(\sigma)$ in $\mathcal{T}(S \setminus \Gamma)$, obtained by pinching the geodesic representatives of $\gamma_1, \dots, \gamma_k$, but otherwise leaving the metric unchanged, that is, by retaining the same Fenchel-Nielsen coordinates. Thus we define a homeomorphism

$$\Pi : \text{Thin}_{\epsilon_1}(S, \Gamma) \rightarrow \mathcal{T}(S \setminus \Gamma) \times U_1 \times \dots \times U_k.$$

Endow $\mathcal{T}(S \setminus \Gamma) \times U_1 \times \dots \times U_k$ with the sup metric

$$d_{L_\Gamma} = \sup\{d_{L(S \setminus \Gamma)}, d_{L(A_1)}, \dots, d_{L(A_k)}\}.$$

Theorem 3.4 (product regions for Lipschitz metric). *For any $\sigma, \tau \in \text{Thin}_{\epsilon_1}(S, \Gamma)$, we have*

$$d_L(\sigma, \tau) \stackrel{+}{\asymp} d_{L_\Gamma}(\Pi(\sigma), \Pi(\tau)).$$

The heart of the proof is Proposition 3.6 below.

3.5. Replacing an arc with a loop. Next, we describe a procedure to replace an arc in ζ_Q with a non-trivial, non-peripheral simple closed curve in Q that has comparable length. We assume that Q is not homeomorphic to a pair of pants. Let κ be a simple geodesic arc in Q whose endpoints lie in ∂Q and which is perpendicular to ∂Q . If the two endpoints of κ lie in distinct components C, C' of ∂Q , then the boundary of a regular neighborhood of $\kappa \cup C \cup C'$ in Q consists of a single curve η . Define $\hat{\kappa}$ to be the geodesic representative of η in S . Note that since Q is not a pair of pants, it follows that η is non-peripheral in Q , and in particular, $\hat{\kappa}$ is contained in Q (see Figure 2(a)).

If both endpoints of κ lie in a single component C of ∂Q , then the boundary of a regular neighborhood of $\kappa \cup C$ has two components (see Figure 2(b)). In this case, define $\hat{\kappa}$ to be the curve of greater length between the geodesic representatives in S of the two components. Note that $\hat{\kappa}$ is non-peripheral in Q and in particular, it is contained in Q . Also note that unlike the preceding case, the choice of $\hat{\kappa}$ depends on the geometry of the surface.




FIGURE 2. Construction of $\hat{\kappa}$.


Lemma 3.5. *Suppose Q is a hyperbolic component of an (ϵ_0, ϵ_1) -decomposition of σ . Let κ be an arc in Q perpendicular to ∂Q and let $\hat{\kappa}$ be the associated simple closed curve constructed above. If $l_\sigma(\hat{\kappa}) > c_0$ for the Margulis constant c_0 , then*

$$l_\sigma(\kappa) \asymp l_\sigma(\hat{\kappa}).$$

Proof. Let C, C' denote the components of ∂Q that contain the endpoints of κ , where we take $C = C'$ if the endpoints lie on the same component. Let γ, γ' denote the geodesic representatives of C, C' in S . By hypothesis, γ and γ' have embedded collars in S , whose boundary components each have length ϵ_0 . Cut the collars in half along γ, γ' and let $\overline{Q}$ be the surface obtained by attaching the half collars around γ, γ' to Q , along C, C' , respectively. (In the case that $C \neq C'$ but $\gamma = \gamma'$ in S , we attach a half-collar around γ to each of C and C' .) Since κ intersects ∂Q perpendicularly, it has a natural extension to a (smooth) geodesic arc $\bar{\kappa}$ with endpoints in $\partial \overline{Q}$ and perpendicular to $\partial \overline{Q}$, as depicted in Figure 2.

First, consider the case when $C \neq C'$. Let P be the pair of pants with boundary components $\gamma, \gamma', \hat{\kappa}$ and consider one of the right-angled hexagons of P , as in Figure 3(a).

Let $a = l(\gamma)/2, a' = l(\gamma')/2$ and let d, d' be the widths of the half-collars around γ, γ' , respectively. Let $b = l(\kappa)$ and $c = l(\hat{\kappa})/2$. By the

FIGURE 3. A hexagon and pentagon of P .

formula for right-angled hexagons, we have

$$(18) \quad \cosh c + \cosh a \cosh a' = \sinh a \sinh a' \cosh(b + d + d').$$

Since $a, a' < \epsilon_1/2$ and since ϵ_1 is smaller than the Margulis constant $c_0 = 0.2629 \dots$ [12], we have that $\sinh a < 2a$ and $\sinh a' < 2a'$. Also, by a straightforward calculation in $\mathbb{H}^2$, we have $\epsilon_0 = a \cosh d = a' \cosh d'$. Therefore, the right-hand side of Equation (18) satisfies

$$\begin{aligned} \sinh a \sinh a' \cosh(b + d + d') &> a \cdot a' \frac{e^{b+d+d'}}{2} \\ &> a \cdot a' \frac{\cosh d \cosh d'}{8} e^b > \frac{\epsilon_0^2 e^b}{8}, \\ \sinh a \sinh a' \cosh(b + d + d') &< 4a \cdot a' \cdot e^{b+d+d'} \\ &< a \cdot a' \cdot \cosh d \cosh d' e^b < \epsilon_0^2 e^b. \end{aligned}$$

On the other hand, since $a, a' < \epsilon_1/2 < \epsilon_0/4 < c_0/4$ and $c > c_0/2$, we have

$$\cosh a \cosh a' < \cosh(a + a') < \cosh \frac{c_0}{2} < \cosh c.$$

Therefore, Equation (18) combined with the three equations above gives

$$\epsilon_0^2 e^b / 16 < \cosh c < \epsilon_0^2 e^b.$$

Hence,

$$|c - b| = \left| \frac{l(\hat{\kappa})}{2} - l(\kappa) \right| < 2 \log \frac{1}{\epsilon_0} + k$$

for some universal constant $k (= 4 \log 2)$. Thus, if $l(\kappa)$ is sufficiently large, the additive error can be absorbed into multiplicative constants to conclude $l(\hat{\kappa}) \asymp l(\kappa)$. If $l(\kappa)$ is not sufficiently large, then $l(\hat{\kappa}) \asymp l(\kappa)$ holds almost tautologically, because $l(\hat{\kappa})$ is bounded above by $2l(\kappa) + 2\epsilon_0$ and is bounded below, by assumption.

Next consider the case where $C = C'$. Let P be the geodesic pair of pants in S filled by $\bar{\kappa} \cup \gamma$. The arc $\bar{\kappa}$ divides the two right-angled hexagons of P into four right-angled pentagons. It is easy to see that the two pentagons that have edges originally contained in $\hat{\kappa}$ are isometric to one another. Let X be either one of them, as in Figure 3(b). Let $b = l(\kappa)/2$, $c = l(\hat{\kappa})/2$ and let d be the width of the half-collar around γ . Let a be the length of the edge of X coming from γ . Now, by the formula for right-angled pentagons, we have

$$\cosh c = \sinh(b + d) \sinh a.$$

It is clear that $a \leq l(\gamma)/2$ and by applying the pentagon formula to the pentagon which together with X makes up a hexagon of P , we see that our choice of $\hat{\kappa}$ implies $a \geq l(\gamma)/4$. Furthermore, as before we have $l(\gamma) \cdot \cosh d = \epsilon_0$ and since $l(\gamma) \leq \epsilon_1$, the assumption that $\epsilon_0/\epsilon_1 > 2$ is sufficient to guarantee that d is large enough that $e^{b+d}/4 < \sinh(b + d)$ holds. And, as above, ϵ_1 is small enough that $a < \sinh a < 2a$. Therefore, we have

$$\begin{aligned} \cosh c &= \sinh(b + d) \sinh a > \frac{e^b e^d}{4} a > e^b \frac{\cosh d}{4} \cdot \frac{l(\gamma)}{4} > \frac{e^b \epsilon_0}{16}. \\ \cosh c &= \sinh(b + d) \sinh a < e^b e^d a < e^b \cdot 2 \cosh d \cdot l(\gamma)/2 = e^b \epsilon_0. \end{aligned}$$

Hence,

$$|c - b| = \left| \frac{l(\hat{\kappa})}{2} - \frac{l(\kappa)}{2} \right| < \log \frac{1}{\epsilon_0} + k$$

for some universal constant $k (= 4 \log 2)$. Thus we conclude as before that $l(\hat{\kappa}) \asymp l(\kappa)$. $\square$

We remark that in the second case above, had we not chosen $\hat{\kappa}$ to be the longer of the two components of $\partial P - \gamma$, then the lemma would not be true. This can be easily seen by considering the construction in reverse as follows. Take a closed curve α in Q of moderate length and a very long arc β with one endpoint on α and the other on a component C of ∂Q . Construct a new arc κ with both endpoints on C by replacing β with two copies of itself very close together and by connecting their two endpoints on α by the longer arc along α . It is easy to see that the pair of pants filled by $\kappa \cup C$ has α as a boundary component, yet, $l(\alpha)/l(\kappa)$ can be made arbitrarily small.

3.6. Proof of product region theorem for Lipschitz metric. For any surface Σ , let $\mathcal{C}(\Sigma)$ be the set of homotopy classes of non-peripheral, non-trivial simple closed curves in Σ . We are now ready to prove:

Proposition 3.6. *Suppose $\mathcal{P}$ is a partial (ϵ_0, ϵ_1) -decomposition for both $\sigma, \tau \in \mathcal{T}(S)$. Then*

$$(19) \quad \sup_{\zeta \in \mathcal{C}(S)} \frac{l_\tau(\zeta)}{l_\sigma(\zeta)} \asymp \max_{Q, A \in \mathcal{P}} \left\{ \sup_{\alpha \in \mathcal{C}(Q)} \frac{l_\tau(\alpha)}{l_\sigma(\alpha)}, \sup_{\beta \in \mathcal{C}(A, \partial A)} \frac{l_\tau(\beta)}{l_\sigma(\beta)} \right\}.$$

Moreover, when taking the maximum, we may assume Q is never a pair of pants.

Proof. By Corollary 3.2, for any curve $\zeta \in \mathcal{C}(S)$ and any $\rho \in \mathcal{T}(S)$ which has $\mathcal{P}$ as a partial (ϵ_0, ϵ_1) -decomposition, we have

$$l_\rho(\zeta) \asymp \sum_{Q, A \in \mathcal{P}} [l_\rho(\zeta_Q) + l_\rho(\zeta_A)],$$

where $\zeta \cap A$ is written ζ_A for short. Applied to σ, τ , this gives

$$(20) \quad \frac{l_\tau(\zeta)}{l_\sigma(\zeta)} \asymp \frac{\sum_{Q, A \in \mathcal{P}} [l_\tau(\zeta_Q) + l_\tau(\zeta_A)]}{\sum_{Q, A \in \mathcal{P}} [l_\sigma(\zeta_Q) + l_\sigma(\zeta_A)]} \leq \max_{Q, A \in \mathcal{P}} \left\{ \frac{l_\tau(\zeta_Q)}{l_\sigma(\zeta_Q)}, \frac{l_\tau(\zeta_A)}{l_\sigma(\zeta_A)} \right\}.$$

Fix Q and write $\zeta_Q = \sum_i m_i \kappa_i + \sum_j n_j \lambda_j$, where κ_i are arcs with endpoints on ∂Q and λ_j are non-peripheral simple closed curves contained in Q . Then

$$(21) \quad \frac{l_\tau(\zeta_Q)}{l_\sigma(\zeta_Q)} \asymp \frac{\sum_i m_i l_\tau(\kappa_i) + \sum_j n_j l_\tau(\lambda_j)}{\sum_i m_i l_\sigma(\kappa_i) + \sum_j n_j l_\sigma(\lambda_j)} \leq \max_{i, j} \left\{ \frac{l_\tau(\kappa_i)}{l_\sigma(\kappa_i)}, \frac{l_\tau(\lambda_j)}{l_\sigma(\lambda_j)} \right\}.$$

The idea is to show that for every i ,

$$(22) \quad \frac{l_\tau(\kappa_i)}{l_\sigma(\kappa_i)} \prec \sup_{\alpha \in \mathcal{C}(Q)} \frac{l_\tau(\alpha)}{l_\sigma(\alpha)},$$

by replacing $\kappa = \kappa_i$ with the associated simple closed curve $\hat{\kappa} = \hat{\kappa}_i$ in Q , as described above. In the case that Q is a pair of pants, it is not hard to see that there are multiplicative constants depending only on ϵ_0 such that $l(\kappa) \asymp i(\kappa, \partial Q)$ and so

$$\frac{l_\tau(\kappa)}{l_\sigma(\kappa)} \asymp \frac{i(\kappa, \partial Q)}{i(\kappa, \partial Q)} \asymp 1.$$

Therefore, it is sufficient to prove Equation (22) assuming that Q is not a pair of pants, so that we may apply Lemma 3.5.

Recall, that in the case the two endpoints of κ lie in the same component of ∂Q , the choice of $\hat{\kappa}$ depends on the geometry of the surface. Let $\hat{\kappa}(\tau), \hat{\kappa}(\sigma)$ denote the curves associated to κ for the two metrics τ, σ , respectively. Note that by definition of $\hat{\kappa}$,

$$l_\sigma(\hat{\kappa}(\tau)) \leq l_\sigma(\hat{\kappa}(\sigma)).$$

Now, if $l_\tau(\hat{\kappa}(\tau)) > c_0$, then applying Lemma 3.5 and using the fact that $l(\hat{\kappa}) \leq 2l(\kappa) + 2\epsilon_0$ always holds, we have

$$\frac{l_\tau(\kappa)}{l_\sigma(\kappa)} \dot{\prec} \frac{l_\tau(\hat{\kappa}(\tau))}{l_\sigma(\kappa)} \dot{\prec} \frac{l_\tau(\hat{\kappa}(\tau))}{l_\sigma(\hat{\kappa}(\sigma))} \leq \frac{l_\tau(\hat{\kappa}(\tau))}{l_\sigma(\hat{\kappa}(\tau))} \leq \sup_{\alpha \in \mathcal{C}(Q)} \frac{l_\tau(\alpha)}{l_\sigma(\alpha)}.$$

If $l_\tau(\hat{\kappa}(\tau)) \leq c_0$, then in the τ -metric, the three boundary curves of the geodesic pair of pants P spanned by $\bar{\kappa} \cup \gamma \cup \gamma'$ (see Lemma 3.5 above) all have length shorter than c_0 . By using the formulas for right-angled pentagons and hexagons as in the proof of Lemma 3.5, it is easy to show that this implies $l_\tau(\kappa)$ is bounded above. Furthermore, since κ meets ∂Q and ∂Q has an embedded regular neighborhood of some definite width depending on ϵ_0 , it follows that $l_\sigma(\kappa)$ is bounded below. Hence,

$$\frac{l_\tau(\kappa)}{l_\sigma(\kappa)} \dot{\prec} \frac{1}{l_\sigma(\kappa)} \dot{\prec} 1.$$

Since the ratio $l_\tau(\kappa)/l_\sigma(\kappa)$ is bounded above, Equation (22) is tautologically satisfied in this case. Thus Equation (22) is proved.

Combined with Equations (20) and (21) we now have

$$\frac{l_\tau(\zeta)}{l_\sigma(\zeta)} \dot{\prec} \max_{Q, A \in \mathcal{P}} \left\{ \sup_{\alpha \in \mathcal{C}(Q)} \frac{l_\tau(\alpha)}{l_\sigma(\alpha)}, \sup_{\beta \in \mathcal{C}(A, \partial A)} \frac{l_\tau(\beta)}{l_\sigma(\beta)} \right\},$$

where by $l_\sigma(\beta), l_\tau(\beta)$ for $\beta \in \mathcal{C}(A, \partial A)$, we mean the length of β in the metrics $\Pi_A(\sigma), \Pi_A(\tau) \in U_{\epsilon_1}(A)$, respectively, as defined in Section 3.4. Therefore, the supremum of the left hand side, taken over all $\zeta \in \mathcal{C}(S)$, is bounded by the quantity on the right hand side.

Finally, since $\mathcal{C}(Q) \subset \mathcal{C}(S)$, it is clear that for every $Q \in \mathcal{P}$

$$\sup_{\zeta \in \mathcal{C}(S)} \frac{l_\tau(\zeta)}{l_\sigma(\zeta)} \geq \max_{Q \in \mathcal{P}} \left\{ \sup_{\alpha \in \mathcal{C}(Q)} \frac{l_\tau(\alpha)}{l_\sigma(\alpha)} \right\}.$$

To complete the proof, we will show that there is a simple closed curve ζ such that

$$\frac{l_\tau(\zeta)}{l_\sigma(\zeta)} \dot{\prec} \sup_{\beta \in \mathcal{C}(A, \partial A)} \frac{l_\tau(\beta)}{l_\sigma(\beta)}.$$

Given an annulus $A \in \mathcal{P}$, suppose that β is a geodesic arc in A with endpoints on ∂A that realizes the supremum on the right. Let α denote the core geodesic of A . If α does not separate S , we can always find a non-trivial arc δ contained in $S \setminus A$ joining the endpoints of β whose length in σ is bounded above by some constant ℓ depending only on ϵ_0 , and such that $\beta \cup \delta$ forms a non-trivial simple closed curve. Let ζ be the

closed curve in the isotopy class of $\beta \cup \delta$. Then we have $l_\rho(\zeta) \asymp l_\rho(\beta)$ and it follows from Corollary 3.2 that $l_\tau(\zeta) \asymp l_\tau(\beta)$. Therefore,

$$\frac{l_\tau(\zeta)}{l_\sigma(\zeta)} \asymp \frac{l_\tau(\beta)}{l_\sigma(\beta)}.$$

In the case that α separates S , we take an additional arc β' in $(A, \partial A)$ disjoint from β . Construct a simple closed curve ζ by joining the pairs of endpoints of β, β' which lie in a common component of ∂A , by arcs δ, δ' in $S \setminus A$, whose lengths in σ are uniformly bounded above. By the same argument as before, we can again show that

$$\frac{l_\tau(\zeta)}{l_\sigma(\zeta)} \asymp \frac{l_\tau(\beta)}{l_\sigma(\beta)}.$$

□

We conclude this section with the proof of Theorem 3.4:

Proof of Theorem 3.4. By Proposition 3.6, we have that

$$d_L(\sigma, \tau) \asymp \max_{Q, A \in \mathcal{P}} \left\{ \log \sup_{\alpha \in \mathcal{C}(Q)} \frac{l_\tau(\alpha)}{l_\sigma(\alpha)}, \log \sup_{\alpha \in \mathcal{C}(Q)} \frac{l_\sigma(\alpha)}{l_\tau(\alpha)}, d_{L(A)}(\sigma, \tau) \right\}.$$

Therefore, to complete the proof, it would be sufficient to show that

$$\sup_{\alpha \in \mathcal{C}(Q)} \frac{l_\tau(\alpha)}{l_\sigma(\alpha)} \asymp \sup_{\alpha \in \mathcal{C}(Q)} \frac{l_{\Pi_{S \setminus \Gamma}(\tau)}(\alpha)}{l_{\Pi_{S \setminus \Gamma}(\sigma)}(\alpha)}.$$

However, it was already shown in [5] that for $\rho \in \text{Thin}_\epsilon(S, \Gamma)$, the space (Q, ρ) embeds K -quasiconformally (in fact, biLipschitz), with uniform K , in $(Q, \pi_{S \setminus \Gamma}(\rho))$. Thus, the lengths of curves in the two spaces are comparable and the theorem follows. □

4. COMPARISON ON A THIN REGION

We now provide an example that illustrates the discrepancy between the Lipschitz and Teichmüller distances stated in the introduction:

Theorem A. *There are sequences $\sigma_n, \tau_n \in \mathcal{T}(S)$ such that, as $n \rightarrow \infty$,*

$$d_L(\sigma_n, \tau_n) \rightarrow 0, \quad d_T(\sigma_n, \tau_n) \rightarrow \infty.$$

Proof. Let σ_n be a hyperbolic metric on S such that there is exactly one short curve γ of length $l_{\sigma_n}(\gamma) = \epsilon_n$ and let $\tau_n = D_\gamma^{T_n}(\sigma_n)$ be the metric obtained from σ_n by T_n Dehn twists around γ . In this case, $l_{\sigma_n}(\gamma) = l_{\tau_n}(\gamma) = \epsilon_n$. Set $\epsilon_n = e^{-P_n}$, $T_n = e^{P_n + q_n}$ and choose the sequences of positive integers P_n, q_n so that

$$P_n \rightarrow \infty, \quad q_n \rightarrow \infty \quad \text{and} \quad \frac{e^{q_n}}{P_n} \rightarrow 0 \quad \text{as} \quad n \rightarrow \infty.$$

On the one hand, it follows Theorem 1.1 that

$$d_{\mathcal{T}}(\sigma_n, \tau_n) \stackrel{+}{\asymp} \log[T_n \epsilon_n] = q_n \rightarrow \infty.$$

On the other hand, it follows from Proposition 3.1 that for a simple closed curve ζ in S , we have

$$\frac{l_{\tau_n}(\zeta)}{l_{\sigma_n}(\zeta)} = \frac{l_{\tau_n}(\zeta_Q) + [\log[\epsilon_0/\epsilon_n] + \epsilon_n \cdot Tw_{\tau_n}(\zeta, \gamma)/2 + O(1)] \cdot i(\zeta, \gamma)}{l_{\sigma_n}(\zeta_Q) + [\log[\epsilon_0/\epsilon_n] + \epsilon_n \cdot Tw_{\sigma_n}(\zeta, \gamma)/2 + O(1)] \cdot i(\zeta, \gamma)},$$

where $O(1)$ represents an error that is independent of ζ, σ_n, τ_n and that is bounded in absolute value by some uniform constant. Since σ_n, τ_n coincide outside of A , we have $l_{\tau_n}(\zeta_Q) = l_{\sigma_n}(\zeta_Q)$. Therefore,

$$\sup_{\zeta} \frac{l_{\tau_n}(\zeta)}{l_{\sigma_n}(\zeta)} \leq \max \left\{ 1, \sup_{\zeta} \frac{2 \log[\epsilon_0/\epsilon_n] + \epsilon_n \cdot Tw_{\tau_n}(\zeta, \gamma) + O(1)}{2 \log[\epsilon_0/\epsilon_n] + \epsilon_n \cdot Tw_{\sigma_n}(\zeta, \gamma) + O(1)} \right\}$$

and by the same reasoning used to deduce Equation (17), the supremum on the right-hand side is equal to

$$\frac{2 \log[1/\epsilon_n] + \epsilon_n \cdot T_n + O(1)}{2 \log[1/\epsilon_n] + O(1)} = \frac{2P_n + e^{q_n} + O(1)}{2P_n + O(1)}.$$

Thus, we have

$$\lim_{n \rightarrow \infty} d_L(\sigma_n, \tau_n) = \lim_{n \rightarrow \infty} \log \frac{2P_n + e^{q_n} + O(1)}{2P_n + O(1)} = 0. \quad \square$$

So far, we have seen that if $\sigma, \tau \in \mathcal{T}(S)$ are both in the thick part then $d_L(\sigma, \tau) \asymp d_{\mathcal{T}}(\sigma, \tau)$, but that if σ, τ have a short curve in common, then the two distances are no longer comparable. The following proposition shows that, in some sense, this is the only way for the distances to diverge.

Proposition 4.1. *If $\sigma, \tau \in \mathcal{T}(S)$ have no short curves in common, then $d_L(\sigma, \tau) \asymp d_{\mathcal{T}}(\sigma, \tau)$.*

Proof. Let Γ_{σ} be the set of curves whose length is less than ϵ_1 at σ and let $\bar{\sigma}$ be the point in the thick part of $\mathcal{T}(S)$ obtained from σ by increasing the length of each curve in Γ_{σ} to ϵ_1 but otherwise leaving the metric unchanged. This, as usual, can be achieved by choosing a marking μ_{σ} of S that contains Γ_{σ} in its pants system and altering the associated Fenchel-Nielsen length coordinates as desired. We define $\bar{\tau}$ analogously by increasing the length of every short curve of τ to ϵ_1 . It follows from Theorem 3.4 and Lemma 3.3 that

$$d_L(\sigma, \bar{\sigma}) \asymp \log \max_{\alpha \in \Gamma_{\sigma}} \left\{ \frac{l_{\bar{\sigma}}(\alpha)}{l_{\sigma}(\alpha)} \right\}, \quad d_L(\tau, \bar{\tau}) \asymp \log \max_{\alpha \in \Gamma_{\tau}} \left\{ \frac{l_{\bar{\tau}}(\alpha)}{l_{\tau}(\alpha)} \right\}.$$

Since curves that are short in σ are not short in τ and vice versa, the above equation implies that

$$(23) \quad d_L(\sigma, \bar{\sigma}) \prec d_L(\sigma, \tau) \text{ and } d_L(\tau, \bar{\tau}) \prec d_L(\sigma, \tau).$$

By the triangle inequality, we also have

$$(24) \quad \begin{aligned} d_L(\sigma, \tau) &\geq d_L(\bar{\sigma}, \bar{\tau}) - d_L(\sigma, \bar{\sigma}) - d_L(\bar{\tau}, \tau), \\ d_L(\sigma, \tau) &\leq d_L(\bar{\sigma}, \bar{\tau}) + d_L(\sigma, \bar{\sigma}) + d_L(\bar{\tau}, \tau) \end{aligned}$$

Combining Equations (23) and (24), we get

$$(25) \quad d_L(\sigma, \tau) \asymp d_L(\bar{\sigma}, \sigma) + d_L(\bar{\sigma}, \bar{\tau}) + d_L(\tau, \bar{\tau}).$$

Analogously, it follows from Theorem 1.1 and Equation (1) that

$$d_T(\sigma, \bar{\sigma}) \prec d_T(\sigma, \tau) \text{ and } d_T(\tau, \bar{\tau}) \prec d_T(\sigma, \tau)$$

and combined with the triangle inequality again, we get

$$(26) \quad d_T(\sigma, \tau) \asymp d_T(\bar{\sigma}, \sigma) + d_T(\bar{\sigma}, \bar{\tau}) + d_T(\tau, \bar{\tau}).$$

Now, by Theorem 3.4, Lemma 3.3, and Theorem 1.1 we have

$$d_L(\sigma, \bar{\sigma}) \asymp d_T(\sigma, \bar{\sigma}) \text{ and } d_L(\tau, \bar{\tau}) \asymp d_T(\tau, \bar{\tau})$$

and by Theorem B we have

$$d_L(\bar{\sigma}, \bar{\tau}) \asymp d_T(\bar{\sigma}, \bar{\tau}).$$

Thus it follows from Equations (25) and (26) that $d_L(\sigma, \tau) \asymp d_T(\sigma, \tau)$, as claimed. $\square$

REFERENCES

- [1] A. Belkhirat, A. Papadopoulos, and M. Troyanov, “Thurston’s weak metric on the Teichmüller space of the torus”, *Trans. Amer. Math. Soc.* **357** (2005) 3311-3324.
- [2] Y. Choi, K. Rafi and C. Series, “Lines of minima are Teichmüller quasi-geodesics”, in preparation.
- [3] S. Kerckhoff, “The asymptotic geometry of Teichmüller space”, *Topology* **19** (1980) 23-41.
- [4] H. Masur and Y. Minsky, “Geometry of the complex of curves II: Hierarchical structure”, *Geom. Funct. Anal.* **10** (2000) 902-974.
- [5] Y. Minsky, “Extremal length estimates and product regions in Teichmüller space”, *Duke Math. J.* **83** (1996) 249-286.
- [6] Y. Minsky, “Teichmüller geodesics and ends of hyperbolic 3-manifolds”, *Topology* **32** (1993) 625-647.
- [7] K. Rafi, “A characterization of short curves of a Teichmüller geodesic”, *Geometry and Topology* **9** (2005) 179-202.
- [8] K. Rafi, “A combinatorial model for the Teichmüller metric”, e-print arXiv:math.GT/0509584.
- [9] K. Rafi, “A thick-thin decomposition for quadratic differentials”, preprint.

- [10] W. Thurston, “Minimal stretch maps between hyperbolic surfaces”, e-print arXiv:math.GT/9801039.
- [11] S. Wolpert, “The length spectra as moduli for compact Riemann surfaces”, *Ann. of Math.* **109** (1979) 323-351.
- [12] A. Yamada, “On Marden’s universal constant of Fuchsian groups”, *Kodai J.* **4** (1981) 266-277.

E-mail address: choiye@math.ucdavis.edu

E-mail address: rafi@math.uconn.edu

URL: <http://www.math.uconn.edu/~rafi/>